12

The Cell Cycle



▲ **Figure 12.1 How do a cell's chromosomes change during cell division?**

KEY CONCEPTS

12.1 Most cell division results in genetically identical daughter cells

12.2 The mitotic phase alternates with interphase in the cell cycle

12.3 The eukaryotic cell cycle is regulated by a molecular control system

OVERVIEW

The Key Roles of Cell Division

The ability of organisms to produce more of their own kind is the one characteristic that best distinguishes living things from nonliving matter. This unique capacity to procreate, like all biological functions, has a cellular basis. Rudolf Virchow, a German physician, put it this way in 1855: "Where a cell exists, there must have been a preexisting cell, just as the animal arises only from an animal and the plant only from a plant." He summarized this concept with the Latin axiom "*Omnis cellula e cellula*," meaning "Every cell from a cell." The continuity of life is based on the reproduction of cells, or **cell division**. The series of fluorescence micrographs in **Figure 12.1** follows an animal cell's chromosomes, from lower left to lower right, as one cell divides into two.

Cell division plays several important roles in life. The division of one prokaryotic cell reproduces an entire organism. The same is true of a unicellular eukaryote **(Figure 12.2a)**. Cell division also enables multicellular eukaryotes to develop from a single cell, like the fertilized egg that gave rise to the two-celled embryo in **Figure 12.2b**. And after such an organism is fully grown, cell division continues to function in renewal and repair, replacing cells that die from normal wear and tear or accidents. For example, dividing cells in your bone marrow continuously make new blood cells **(Figure 12.2c)**.

The cell division process is an integral part of the **cell cycle**, the life of a cell from the time it is first formed from a dividing parent cell until its own division into two daughter cells. (Our use of the words *daughter* or *sister* in relation to cells is not meant to imply gender.) Passing identical genetic material to cellular offspring is a crucial function of cell division. In this chapter, you will learn how this process occurs. After studying the cellular mechanics of cell division in eukaryotes and bacteria, you will learn about the molecular control system that regulates progress through the eukaryotic cell cycle and what happens when the control system malfunctions. Because a breakdown in cell cycle control plays a major role in cancer development, this aspect of cell biology is an active area of research.



◀ **(a) Reproduction.** An amoeba, a single-celled eukaryote, is dividing into two cells. Each new cell will be an individual organism (LM).



▶ **(b) Growth and development.** This micrograph shows a sand dollar embryo shortly after the fertilized egg divided, forming two cells (LM).



◀ **(c) Tissue renewal.** These dividing bone marrow cells will give rise to new blood cells (LM).

▲ **Figure 12.2 The functions of cell division.**

CONCEPT 12.1

Most cell division results in genetically identical daughter cells

The reproduction of an assembly as complex as a cell cannot occur by a mere pinching in half; a cell is not like a soap bubble that simply enlarges and splits in two. In both prokaryotes and eukaryotes, most cell division involves the distribution of identical genetic material—DNA—to two daughter cells. (The exception is meiosis, the special type of eukaryotic cell division that can produce sperm and eggs.) What is most remarkable about cell division is the fidelity with which the DNA is passed along from one generation of cells to the next. A dividing cell duplicates its DNA, allocates the two copies to opposite ends of the cell, and only then splits into daughter cells. After we describe the distribution of DNA during cell division in animal and plant cells, we'll consider the process in other eukaryotes as well as in bacteria.

Cellular Organization of the Genetic Material

A cell's endowment of DNA, its genetic information, is called its **genome**. Although a prokaryotic genome is often a single DNA molecule, eukaryotic genomes usually consist of a number of DNA molecules. The overall length of DNA in a eukaryotic cell is enormous. A typical human cell, for example, has about 2 m of DNA—a length about 250,000 times greater than the cell's diameter. Yet before the cell can divide to form genetically identical daughter cells, all of this DNA must be copied, or replicated, and then the two copies must be separated so that each daughter cell ends up with a complete genome.

The replication and distribution of so much DNA is manageable because the DNA molecules are packaged into structures called **chromosomes**, so named because they take up certain dyes used in microscopy (from the Greek *chroma*, color, and *soma*, body) **(Figure 12.3)**. Each eukaryotic chromosome consists of one very long, linear DNA molecule associated with many proteins (see Figure 6.9). The DNA molecule carries several hundred to a few thousand genes, the units of information that specify an organism's inherited traits. The associated proteins maintain the structure of the chromosome and help control the activity of the genes. Together, the entire complex of DNA and proteins that is the building material of chromosomes is referred to as **chromatin**. As you will soon see, the chromatin of a chromosome varies in its degree of condensation during the process of cell division.

Every eukaryotic species has a characteristic number of chromosomes in each cell nucleus. For example, the nuclei of human **somatic cells** (all body cells except the reproductive cells) each contain 46 chromosomes, made up of two sets of 23, one set inherited from each parent. Reproductive cells, or **gametes**—sperm and eggs—have half as many chromosomes as somatic cells, or one set of 23 chromosomes in humans. The number of chromosomes in somatic cells varies widely among species: 18 in cabbage plants, 48 in chimpanzees, 56 in elephants, 90 in hedgehogs, and 148 in one species of alga. We'll now consider how these chromosomes behave during cell division.



▲ **Figure 12.3 Eukaryotic chromosomes.** Chromosomes (stained purple) are visible within the nucleus of this cell from an African blood lily. The thinner red threads in the surrounding cytoplasm are the cytoskeleton. The cell is preparing to divide (LM).

Distribution of Chromosomes During Eukaryotic Cell Division

When a cell is not dividing, and even as it replicates its DNA in preparation for cell division, each chromosome is in the form of a long, thin chromatin fiber. After DNA replication, however, the chromosomes condense as a part of cell division: Each chromatin fiber becomes densely coiled and folded, making the chromosomes much shorter and so thick that we can see them with a light microscope.

Each duplicated chromosome has two **sister chromatids**, which are joined copies of the original chromosome **(Figure 12.4)**. The two chromatids, each containing an identical DNA molecule, are initially attached all along their lengths by protein complexes called *cohesins*; this attachment is known as *sister chromatid cohesion*. Each sister chromatid has a **centromere**, a region containing specific DNA sequences



▲ **Figure 12.4 A highly condensed, duplicated human chromosome (SEM).**

DRAW IT *Circle one sister chromatid of the chromosome in this micrograph.*



▶ Figure 12.5 Chromosome duplication and distribution during cell division.

? *How many chromatid arms does the chromosome in 2 have?*

where the chromatid is attached most closely to its sister chromatid. This attachment is mediated by proteins bound to the centromeric DNA sequences and gives the condensed, duplicated chromosome a narrow "waist." The part of a chromatid on either side of the centromere is referred to as an *arm* of the chromatid. (An uncondensed, unduplicated chromosome has a single centromere and two arms.)

Later in the cell division process, the two sister chromatids of each duplicated chromosome separate and move into two new nuclei, one forming at each end of the cell. Once the sister chromatids separate, they are no longer called sister chromatids but are considered individual chromosomes. Thus, each new nucleus receives a collection of chromosomes identical to that of the parent cell **(Figure 12.5)**. **Mitosis**, the division of the genetic material in the nucleus, is usually followed immediately by **cytokinesis**, the division of the cytoplasm. One cell has become two, each the genetic equivalent of the parent cell.

What happens to the chromosome number as we follow the human life cycle through the generations? You inherited 46 chromosomes, one set of 23 from each parent. They were combined in the nucleus of a single cell when a sperm from your father united with an egg from your mother, forming a fertilized egg, or zygote. Mitosis and cytokinesis produced the 200 trillion somatic cells that now make up your body, and the same processes continue to generate new cells to replace dead and damaged ones. In contrast, you produce gametes—eggs or sperm—by a variation of cell division called *meiosis*, which yields nonidentical daughter cells that have only one set of chromosomes, half as many chromosomes as the parent cell. Meiosis in humans occurs only in the gonads (ovaries or testes). In each generation, meiosis reduces the chromosome number from 46 (two sets of chromosomes) to 23 (one set). Fertilization fuses two gametes together and returns the chromosome number to 46, and mitosis conserves that number in every somatic cell nucleus of the new individual. In Chapter 13, we will examine the role of meiosis in reproduction and inheritance in more detail. In the remainder of this chapter, we focus on mitosis and the rest of the cell cycle in eukaryotes.

CONCEPT CHECK 12.1

1. How many chromatids are in a duplicated chromosome?
2. **WHAT IF?** A chicken has 78 chromosomes in its somatic cells. How many chromosomes did the chicken inherit from each parent? How many chromosomes are in each of the chicken's gametes? How many chromosomes will be in each somatic cell of the chicken's offspring?

For suggested answers, see Appendix A.

CONCEPT 12.2

The mitotic phase alternates with interphase in the cell cycle

In 1882, a German anatomist named Walther Flemming developed dyes that allowed him to observe, for the first time, the behavior of chromosomes during mitosis and cytokinesis. (In fact, Flemming coined the terms *mitosis* and *chromatin*.) During the period between one cell division and the next, it appeared to Flemming that the cell was simply growing larger. But we now know that many critical events occur during this stage in the life of a cell.

Phases of the Cell Cycle

Mitosis is just one part of the cell cycle **(Figure 12.6)**. In fact, the **mitotic (M) phase**, which includes both mitosis and cytokinesis, is usually the shortest part of the cell cycle. Mitotic cell division alternates with a much longer stage called **interphase**, which often accounts for about 90% of the cycle. During interphase, a cell that is about to divide grows and copies its chromosomes in preparation for cell division. Interphase can be divided into subphases: the **G_1 phase** ("first gap"), the **S phase** ("synthesis"), and the **G_2 phase** ("second gap"). During all three subphases, a cell that will eventually divide grows by producing proteins and cytoplasmic organelles such as mitochondria and endoplasmic reticulum. However, chromosomes are duplicated only during the S phase. (We will discuss synthesis of DNA in Chapter 16.) Thus, a cell grows (G_1), continues to grow as it copies its chromosomes (S), grows more as it completes preparations for cell division (G_2), and divides (M). The daughter cells may then repeat the cycle.

A particular human cell might undergo one division in 24 hours. Of this time, the M phase would occupy less than 1 hour, while the S phase might occupy about 10–12 hours, or about half the cycle. The rest of the time would be apportioned between the G_1 and G_2 phases. The G_2 phase usually takes 4–6 hours; in our example, G_1 would occupy about 5–6 hours. G_1 is the most variable in length in different types of cells. Some cells in a multicellular organism divide very infrequently or not at all. These cells spend their time in G_1 (or a related phase called G_0) doing their job in the organism—a nerve cell carries impulses, for example.

Mitosis is conventionally broken down into five stages: **prophase**, **prometaphase**, **metaphase**, **anaphase**, and **telophase**. Overlapping with the latter stages of mitosis, cytokinesis completes the mitotic phase. **Figure 12.7**, on the next two pages, describes these stages in an animal cell. Study this figure thoroughly before progressing to the next two sections, which examine mitosis and cytokinesis more closely.



▲ **Figure 12.6 The cell cycle.** In a dividing cell, the mitotic (M) phase alternates with interphase, a growth period. The first part of interphase (G_1) is followed by the S phase, when the chromosomes duplicate; G_2 is the last part of interphase. In the M phase, mitosis distributes the daughter chromosomes to daughter nuclei, and cytokinesis divides the cytoplasm, producing two daughter cells. The relative durations of G_1, S, and G_2 may vary.

The Mitotic Spindle: *A Closer Look*

Many of the events of mitosis depend on the **mitotic spindle**, which begins to form in the cytoplasm during prophase. This structure consists of fibers made of microtubules and associated proteins. While the mitotic spindle assembles, the other microtubules of the cytoskeleton partially disassemble, providing the material used to construct the spindle. The spindle microtubules elongate (polymerize) by incorporating more subunits of the protein tubulin (see Table 6.1) and shorten (depolymerize) by losing subunits.

In animal cells, the assembly of spindle microtubules starts at the **centrosome**, a subcellular region containing material that functions throughout the cell cycle to organize the cell's microtubules. (It is also called the *microtubule-organizing center*.) A pair of centrioles is located at the center of the centrosome, but they are not essential for cell division: If the centrioles are destroyed with a laser microbeam, a spindle nevertheless forms during mitosis. In fact, centrioles are not even present in plant cells, which do form mitotic spindles.

During interphase in animal cells, the single centrosome duplicates, forming two centrosomes, which remain together near the nucleus. The two centrosomes move apart during prophase and prometaphase of mitosis as spindle microtubules grow out from them. By the end of prometaphase, the two centrosomes, one at each pole of the spindle, are at opposite ends of the cell. An **aster**, a radial array of short microtubules, extends from each centrosome. The spindle includes the centrosomes, the spindle microtubules, and the asters.

Each of the two sister chromatids of a duplicated chromosome has a **kinetochore**, a structure of proteins associated with specific sections of chromosomal DNA at each centromere. The chromosome's two kinetochores face in opposite directions. During prometaphase, some of the spindle microtubules attach to the kinetochores; these are called kinetochore microtubules. (The number of microtubules attached to a kinetochore varies among species, from one microtubule in yeast cells to 40 or so in some mammalian cells.) When one of a chromosome's kinetochores is "captured" by microtubules, the chromosome begins to move toward the pole from which those microtubules extend. However, this movement is checked as soon as microtubules from the opposite pole attach to the other kinetochore. What happens next is like a tug-of-war that ends in a draw. The chromosome moves first in one direction, then the other, back and forth, finally settling midway between the two ends of the cell. At metaphase, the centromeres of all the duplicated chromosomes are on a plane midway between the spindle's

▼ **Figure 12.7**

Exploring Mitosis in an Animal Cell



G_2 of Interphase

- A nuclear envelope encloses the nucleus.
- The nucleus contains one or more nucleoli (singular, *nucleolus*).
- Two centrosomes have formed by duplication of a single centrosome. Centrosomes are regions in animal cells that organize the microtubules of the spindle. Each centrosome contains two centrioles.
- Chromosomes, duplicated during S phase, cannot be seen individually because they have not yet condensed.

The light micrographs show dividing lung cells from a newt, which has 22 chromosomes in its somatic cells. Chromosomes appear blue, microtubules green, and intermediate filaments red. For simplicity, the drawings show only 6 chromosomes.

Prophase

- The chromatin fibers become more tightly coiled, condensing into discrete chromosomes observable with a light microscope.
- The nucleoli disappear.
- Each duplicated chromosome appears as two identical sister chromatids joined at their centromeres and, in some species, all along their arms by cohesins (sister chromatid cohesion).
- The mitotic spindle (named for its shape) begins to form. It is composed of the centrosomes and the microtubules that extend from them. The radial arrays of shorter microtubules that extend from the centrosomes are called asters ("stars").
- The centrosomes move away from each other, propelled partly by the lengthening microtubules between them.

Prometaphase

- The nuclear envelope fragments.
- The microtubules extending from each centrosome can now invade the nuclear area.
- The chromosomes have become even more condensed.
- Each of the two chromatids of each chromosome now has a kinetochore, a specialized protein structure at the centromere.
- Some of the microtubules attach to the kinetochores, becoming "kinetochore microtubules," which jerk the chromosomes back and forth.
- Nonkinetochore microtubules interact with those from the opposite pole of the spindle.

? *How many molecules of DNA are in the prometaphase drawing? How many molecules per chromosome? How many double helices are there per chromosome? Per chromatid?*



Metaphase

- The centrosomes are now at opposite poles of the cell.
- The chromosomes convene at the *metaphase plate*, a plane that is equidistant between the spindle's two poles. The chromosomes' centromeres lie at the metaphase plate.
- For each chromosome, the kinetochores of the sister chromatids are attached to kinetochore microtubules coming from opposite poles.

ANIMATION BioFlix® Visit the Study Area at **www.masteringbiology.com** for the BioFlix® 3-D Animation on Mitosis.

Anaphase

- Anaphase is the shortest stage of mitosis, often lasting only a few minutes.
- Anaphase begins when the cohesin proteins are cleaved. This allows the two sister chromatids of each pair to part suddenly. Each chromatid thus becomes a full-fledged chromosome.
- The two liberated daughter chromosomes begin moving toward opposite ends of the cell as their kinetochore microtubules shorten. Because these microtubules are attached at the centromere region, the chromosomes move centromere first (at about 1 µm/min).
- The cell elongates as the nonkinetochore microtubules lengthen.
- By the end of anaphase, the two ends of the cell have equivalent—and complete—collections of chromosomes.

Telophase

- Two daughter nuclei form in the cell. Nuclear envelopes arise from the fragments of the parent cell's nuclear envelope and other portions of the endomembrane system.
- Nucleoli reappear.
- The chromosomes become less condensed.
- Any remaining spindle microtubules are depolymerized.
- Mitosis, the division of one nucleus into two genetically identical nuclei, is now complete.

Cytokinesis

- The division of the cytoplasm is usually well under way by late telophase, so the two daughter cells appear shortly after the end of mitosis.
- In animal cells, cytokinesis involves the formation of a cleavage furrow, which pinches the cell in two.

two poles. This plane is called the **metaphase plate**, which is an imaginary rather than an actual cellular structure **(Figure 12.8)**. Meanwhile, microtubules that do not attach to kinetochores have been elongating, and by metaphase they overlap and interact with other nonkinetochore microtubules from the opposite pole of the spindle. (These are sometimes called "polar" microtubules.) By metaphase, the microtubules of the asters have also grown and are in contact with the plasma membrane. The spindle is now complete.



▲ **Figure 12.8 The mitotic spindle at metaphase.** The kinetochores of each chromosome's two sister chromatids face in opposite directions. Here, each kinetochore is attached to a *cluster* of kinetochore microtubules extending from the nearest centrosome. Nonkinetochore microtubules overlap at the metaphase plate (TEMs).

DRAW IT *On the lower micrograph, draw a line indicating the position of the metaphase plate. Circle an aster. Draw arrows indicating the directions of chromosome movement once anaphase begins.*

The structure of the completed spindle correlates well with its function during anaphase. Anaphase commences suddenly when the cohesins holding together the sister chromatids of each chromosome are cleaved by an enzyme called *separase*. Once the chromatids become separate, full-fledged chromosomes, they move toward opposite ends of the cell.

How do the kinetochore microtubules function in this poleward movement of chromosomes? Apparently, two mechanisms are in play, both involving motor proteins. (To review how motor proteins move an object along a microtubule, see Figure 6.21.) A clever experiment carried out in 1987 suggested that motor proteins on the kinetochores "walk" the chromosomes along the microtubules, which depolymerize at their kinetochore ends after the motor proteins have passed **(Figure 12.9)**. (This is referred to as the "Pacman" mechanism because of its resemblance to the arcade game character that moves by eating all the dots in its path.) However, other researchers, working with different cell types or cells from other species, have shown that chromosomes are "reeled in" by motor proteins at the spindle poles and that the microtubules depolymerize after they pass by these motor proteins. The general consensus now is that both mechanisms are used and that their relative contributions vary among cell types.

In a dividing animal cell, the nonkinetochore microtubules are responsible for elongating the whole cell during anaphase. Nonkinetochore microtubules from opposite poles overlap each other extensively during metaphase (see Figure 12.8). During anaphase, the region of overlap is reduced as motor proteins attached to the microtubules walk them away from one another, using energy from ATP. As the microtubules push apart from each other, their spindle poles are pushed apart, elongating the cell. At the same time, the microtubules lengthen somewhat by the addition of tubulin subunits to their overlapping ends. As a result, the microtubules continue to overlap.

At the end of anaphase, duplicate groups of chromosomes have arrived at opposite ends of the elongated parent cell. Nuclei re-form during telophase. Cytokinesis generally begins during anaphase or telophase, and the spindle eventually disassembles by depolymerization of microtubules.

Cytokinesis: *A Closer Look*

In animal cells, cytokinesis occurs by a process known as **cleavage**. The first sign of cleavage is the appearance of a **cleavage furrow**, a shallow groove in the cell surface near the old metaphase plate **(Figure 12.10a)**. On the cytoplasmic side of the furrow is a contractile ring of actin microfilaments associated with molecules of the protein myosin. The actin microfilaments interact with the myosin molecules, causing the ring to contract. The contraction of the dividing cell's ring of microfilaments is like the pulling of a drawstring. The cleavage furrow deepens until the parent cell is pinched in two, producing two completely separated cells, each with its own nucleus and share of cytosol, organelles, and other subcellular structures.

▼ Figure 12.9

INQUIRY

At which end do kinetochore microtubules shorten during anaphase?

EXPERIMENT Gary Borisy and colleagues at the University of Wisconsin wanted to determine whether kinetochore microtubules depolymerize at the kinetochore end or the pole end as chromosomes move toward the poles during mitosis. First they labeled the microtubules of a pig kidney cell in early anaphase with a yellow fluorescent dye.



Then they marked a region of the kinetochore microtubules between one spindle pole and the chromosomes by using a laser to eliminate the fluorescence from that region, while leaving the microtubules intact (see below). As anaphase proceeded, they monitored the changes in microtubule length on either side of the mark.



RESULTS As the chromosomes moved poleward, the microtubule segments on the kinetochore side of the mark shortened, while those on the spindle pole side stayed the same length.



CONCLUSION During anaphase in this cell type, chromosome movement is correlated with kinetochore microtubules shortening at their kinetochore ends and not at their spindle pole ends. This experiment supports the hypothesis that during anaphase, a chromosome is walked along a microtubule as the microtubule depolymerizes at its kinetochore end, releasing tubulin subunits.



SOURCE G. J. Gorbsky, P. J. Sammak, and G. G. Borisy, Chromosomes move poleward in anaphase along stationary microtubules that coordinately disassemble from their kinetochore ends, *Journal of Cell Biology* 104:9–18 (1987).

WHAT IF? If this experiment had been done on a cell type in which "reeling in" at the poles was the main cause of chromosome movement, how would the mark have moved relative to the poles? How would the microtubule lengths have changed?

▼ **Figure 12.10 Cytokinesis in animal and plant cells.**

(a) Cleavage of an animal cell (SEM)



(b) Cell plate formation in a plant cell (TEM)



1 Prophase. The chromosomes are condensing and the nucleolus is beginning to disappear. Although not yet visible in the micrograph, the mitotic spindle is starting to form.

2 Prometaphase. Discrete chromosomes are now visible; each consists of two aligned, identical sister chromatids. Later in prometaphase, the nuclear envelope will fragment.

3 Metaphase. The spindle is complete, and the chromosomes, attached to microtubules at their kinetochores, are all at the metaphase plate.

4 Anaphase. The chromatids of each chromosome have separated, and the daughter chromosomes are moving to the ends of the cell as their kinetochore microtubules shorten.

5 Telophase. Daughter nuclei are forming. Meanwhile, cytokinesis has started: The cell plate, which will divide the cytoplasm in two, is growing toward the perimeter of the parent cell.

▲ **Figure 12.11 Mitosis in a plant cell.** These light micrographs show mitosis in cells of an onion root.

Cytokinesis in plant cells, which have cell walls, is markedly different. There is no cleavage furrow. Instead, during telophase, vesicles derived from the Golgi apparatus move along microtubules to the middle of the cell, where they coalesce, producing a **cell plate** (Figure 12.10b). Cell wall materials carried in the vesicles collect in the cell plate as it grows. The cell plate enlarges until its surrounding membrane fuses with the plasma membrane along the perimeter of the cell. Two daughter cells result, each with its own plasma membrane. Meanwhile, a new cell wall arising from the contents of the cell plate has formed between the daughter cells.

Figure 12.11 is a series of micrographs of a dividing plant cell. Examining this figure will help you review mitosis and cytokinesis.

Binary Fission in Bacteria

Prokaryotes (bacteria and archaea) can undergo a type of reproduction in which the cell grows to roughly double its size and then divides to form two cells. The term **binary fission**, meaning "division in half," refers to this process and to the asexual reproduction of single-celled eukaryotes, such as the amoeba in Figure 12.2a. However, the process in eukaryotes involves mitosis, while that in prokaryotes does not.

In bacteria, most genes are carried on a single *bacterial chromosome* that consists of a circular DNA molecule and associated proteins. Although bacteria are smaller and simpler than eukaryotic cells, the challenge of replicating their genomes in an orderly fashion and distributing the copies equally to two daughter cells is still formidable. The chromosome of the bacterium *Escherichia coli*, for example, when it is fully stretched out, is about 500 times as long as the cell. For such a long chromosome to fit within the cell requires that it be highly coiled and folded.

In *E. coli*, the process of cell division is initiated when the DNA of the bacterial chromosome begins to replicate at a specific place on the chromosome called the **origin of replication**, producing two origins. As the chromosome continues to replicate, one origin moves rapidly toward the opposite end of the cell (Figure 12.12). While the chromosome is replicating, the cell elongates. When replication is complete and the bacterium has reached about twice its initial size, its plasma membrane pinches inward, dividing the parent *E. coli* cell into two daughter cells. In this way, each cell inherits a complete genome.

Using the techniques of modern DNA technology to tag the origins of replication with molecules that glow green in fluorescence microscopy (see Figure 6.3), researchers have directly observed the movement of bacterial chromosomes. This movement is reminiscent of the poleward movements of the centromere regions of eukaryotic chromosomes during anaphase of mitosis, but bacteria don't have visible mitotic spindles or even microtubules. In most bacterial species studied, the two origins of replication end up at opposite ends of the cell or in some other very specific location, possibly anchored there by one or more proteins. How bacterial chromosomes move and how their specific location is established and maintained are still not fully understood. However, several proteins have been identified that play important roles: One resembling eukaryotic actin apparently functions in bacterial chromosome movement during cell division, and another that is related to tubulin seems to help pinch the plasma membrane inward, separating the two bacterial daughter cells.



▲ **Figure 12.12 Bacterial cell division by binary fission.** The bacterium *E. coli*, shown here, has a single, circular chromosome.

The Evolution of Mitosis

EVOLUTION Given that prokaryotes preceded eukaryotes on Earth by more than a billion years, we might hypothesize that mitosis evolved from simpler prokaryotic mechanisms of cell reproduction. The fact that some of the proteins involved in bacterial binary fission are related to eukaryotic proteins that function in mitosis supports that hypothesis.

As eukaryotes evolved, along with their larger genomes and nuclear envelopes, the ancestral process of binary fission, seen today in bacteria, somehow gave rise to mitosis. **Figure 12.13** shows some variations on cell division in different groups of organisms. These processes may be similar to mechanisms used by ancestral species and thus may resemble steps in the evolution of mitosis from a binary fission-like process presumably carried out by very early bacteria. Possible intermediate stages are suggested by two unusual types of nuclear division found today in certain unicellular eukaryotes—dinoflagellates, diatoms, and some yeasts. These two modes of nuclear division are thought to be cases where ancestral mechanisms have remained relatively unchanged over evolutionary time. In both types, the nuclear envelope remains intact, in contrast to what happens in most eukaryotic cells.



(a) Bacteria. During binary fission in bacteria, the origins of the daughter chromosomes move to opposite ends of the cell. The mechanism is not fully understood, but proteins may anchor the daughter chromosomes to specific sites on the plasma membrane.



(b) Dinoflagellates. In unicellular protists called dinoflagellates, the chromosomes attach to the nuclear envelope, which remains intact during cell division. Microtubules pass through the nucleus inside cytoplasmic tunnels, reinforcing the spatial orientation of the nucleus, which then divides in a process reminiscent of bacterial binary fission.



(c) Diatoms and some yeasts. In two other groups of unicellular protists, diatoms and some yeasts, the nuclear envelope also remains intact during cell division. In these organisms, the microtubules form a spindle *within* the nucleus. Microtubules separate the chromosomes, and the nucleus splits into two daughter nuclei.



(d) Most eukaryotes. In most other eukaryotes, including plants and animals, the spindle forms outside the nucleus, and the nuclear envelope breaks down during mitosis. Microtubules separate the chromosomes, and two nuclear envelopes then form.

▲ **Figure 12.13 Mechanisms of cell division in several groups of organisms.** Some unicellular eukaryotes existing today have mechanisms of cell division that may resemble intermediate steps in the evolution of mitosis. Except for (a), these schematic diagrams do not show cell walls.

CONCEPT CHECK 12.2

1. How many chromosomes are shown in the diagram in Figure 12.8? Are they duplicated? How many chromatids are shown?
2. Compare cytokinesis in animal cells and plant cells.
3. What is the function of nonkinetochore microtubules?
4. Compare the roles of tubulin and actin during eukaryotic cell division with the roles of tubulin-like and actin-like proteins during bacterial binary fission.
5. **MAKE CONNECTIONS** What other functions do actin and tubulin carry out? Name the proteins they interact with to do so. (Review Figures 6.21a and 6.27a.)
6. **WHAT IF?** During which stages of the cell cycle does a chromosome consist of two identical chromatids?

For suggested answers, see Appendix A.

CONCEPT 12.3

The eukaryotic cell cycle is regulated by a molecular control system

The timing and rate of cell division in different parts of a plant or animal are crucial to normal growth, development, and maintenance. The frequency of cell division varies with the type of cell. For example, human skin cells divide frequently throughout life, whereas liver cells maintain the ability to divide but keep it in reserve until an appropriate need arises—say, to repair a wound. Some of the most specialized cells, such as fully formed nerve cells and muscle cells, do not divide at all in a mature human. These cell cycle differences result from regulation at the molecular level. The mechanisms of this regulation are of intense interest, not only for understanding the life cycles of normal cells but also for understanding how cancer cells manage to escape the usual controls.

Evidence for Cytoplasmic Signals

What controls the cell cycle? One reasonable hypothesis might be that each event in the cell cycle merely leads to the next, as in a simple metabolic pathway. According to this hypothesis, the replication of chromosomes in the S phase, for example, might cause cell growth during the G_2 phase, which might in turn lead inevitably to the onset of mitosis. However, this hypothesis, which proposes a pathway that is not subject to either internal or external regulation, turns out to be incorrect.

In the early 1970s, a variety of experiments led to an alternative hypothesis: that the cell cycle is driven by specific signaling molecules present in the cytoplasm. Some of the first strong evidence for this hypothesis came from experiments with mammalian cells grown in culture. In these experiments, two cells in different phases of the cell cycle were fused to form a single cell with two nuclei. If one of the original cells was in the S phase and the other was in G_1, the G_1 nucleus immediately entered the S phase, as though stimulated by signaling molecules present in the cytoplasm of the first cell. Similarly, if a cell undergoing mitosis (M phase) was fused with another cell in any stage of its cell cycle, even G_1, the second nucleus immediately entered mitosis, with condensation of the chromatin and formation of a mitotic spindle **(Figure 12.14)**.

▼ **Figure 12.14** **INQUIRY**

Do molecular signals in the cytoplasm regulate the cell cycle?

EXPERIMENT Researchers at the University of Colorado wondered whether a cell's progression through the cell cycle is controlled by cytoplasmic molecules. To investigate this, they selected cultured mammalian cells that were at different phases of the cell cycle and induced them to fuse. Two such experiments are shown here.



CONCLUSION The results of fusing a G_1 cell with a cell in the S or M phase of the cell cycle suggest that molecules present in the cytoplasm during the S or M phase control the progression to those phases.

SOURCE R. T. Johnson and P. N. Rao, Mammalian cell fusion: Induction of premature chromosome condensation in interphase nuclei, *Nature* 226:717–722 (1970).

WHAT IF? If the progression of phases did not depend on cytoplasmic molecules and each phase began when the previous one was complete, how would the results have differed?

The Cell Cycle Control System

The experiment shown in Figure 12.14 and other experiments on animal cells and yeasts demonstrated that the sequential events of the cell cycle are directed by a distinct **cell cycle control system**, a cyclically operating set of molecules in the cell that both triggers and coordinates key events



▲ **Figure 12.15 Mechanical analogy for the cell cycle control system.** In this diagram of the cell cycle, the flat "stepping stones" around the perimeter represent sequential events. Like the control device of an automatic washer, the cell cycle control system proceeds on its own, driven by a built-in clock. However, the system is subject to internal and external regulation at various checkpoints, of which three are shown (red).



(a) If a cell receives a go-ahead signal at the G_1 checkpoint, the cell continues on in the cell cycle.

(b) If a cell does not receive a go-ahead signal at the G_1 checkpoint, the cell exits the cell cycle and goes into G_0, a nondividing state.

▲ **Figure 12.16 The G_1 checkpoint.**

WHAT IF? *What might be the result if the cell ignored the checkpoint and progressed through the cell cycle?*

in the cell cycle. The cell cycle control system has been compared to the control device of an automatic washing machine **(Figure 12.15)**. Like the washer's timing device, the cell cycle control system proceeds on its own, according to a built-in clock. However, just as a washer's cycle is subject to both internal control (such as the sensor that detects when the tub is filled with water) and external adjustment (such as activation of the start mechanism), the cell cycle is regulated at certain checkpoints by both internal and external signals.

A **checkpoint** in the cell cycle is a control point where stop and go-ahead signals can regulate the cycle. (The signals are transmitted within the cell by the kinds of signal transduction pathways discussed in Chapter 11.) Animal cells generally have built-in stop signals that halt the cell cycle at checkpoints until overridden by go-ahead signals. Many signals registered at checkpoints come from cellular surveillance mechanisms inside the cell. These signals report whether crucial cellular processes that should have occurred by that point have in fact been completed correctly and thus whether or not the cell cycle should proceed. Checkpoints also register signals from outside the cell, as we will discuss later. Three major checkpoints are found in the G_1, G_2, and M phases (see Figure 12.15).

For many cells, the G_1 checkpoint—dubbed the "restriction point" in mammalian cells—seems to be the most important. If a cell receives a go-ahead signal at the G_1 checkpoint, it will usually complete the G_1, S, G_2, and M phases and divide. If it does not receive a go-ahead signal at that point, it will exit the cycle, switching into a nondividing state called the **G_0 phase** **(Figure 12.16)**. Most cells of the human body are actually in the G_0 phase. As mentioned earlier, mature nerve cells and muscle cells never divide. Other cells, such as liver cells, can be "called back" from the G_0 phase to the cell cycle by external cues, such as growth factors released during injury.

To understand how cell cycle checkpoints work, we first need to see what kinds of molecules make up the cell cycle control system (the molecular basis for the cell cycle clock) and how a cell progresses through the cycle. Then we will consider the internal and external checkpoint signals that can make the clock pause or continue.

The Cell Cycle Clock: Cyclins and Cyclin-Dependent Kinases

Rhythmic fluctuations in the abundance and activity of cell cycle control molecules pace the sequential events of the cell cycle. These regulatory molecules are mainly proteins of two types: protein kinases and cyclins. Protein kinases are enzymes that activate or inactivate other proteins by phosphorylating them (see Chapter 11). Particular protein kinases give the go-ahead signals at the G_1 and G_2 checkpoints.

Many of the kinases that drive the cell cycle are actually present at a constant concentration in the growing cell, but much of the time they are in an inactive form. To be active, such a kinase must be attached to a **cyclin**, a protein that gets its name from its cyclically fluctuating concentration in the cell. Because of this requirement, these kinases are called **cyclin-dependent kinases**, or **Cdks**. The activity of a Cdk rises and falls with changes in the concentration of its cyclin partner. **Figure 12.17a**, on the next page, shows the fluctuating activity of **MPF**, the cyclin-Cdk complex that was discovered first (in frog eggs). Note that the peaks of MPF activity correspond to the peaks of cyclin concentration. The cyclin level rises during the S and G_2 phases and then falls abruptly during M phase.

The initials MPF stand for "maturation-promoting factor," but we can think of MPF as "M-phase-promoting factor" because it triggers the cell's passage past the G_2 checkpoint into



(a) Fluctuation of MPF activity and cyclin concentration during the cell cycle



(b) Molecular mechanisms that help regulate the cell cycle

▲ **Figure 12.17 Molecular control of the cell cycle at the G_2 checkpoint.** The steps of the cell cycle are timed by rhythmic fluctuations in the activity of cyclin-dependent kinases (Cdks). Here we focus on a cyclin-Cdk complex in animal cells called MPF, which acts at the G_2 checkpoint as a go-ahead signal, triggering the events of mitosis.

? *Explain how the events in the diagram in (b) are related to the "Time" axis of the graph in (a).*

M phase **(Figure 12.17b)**. When cyclins that accumulate during G_2 associate with Cdk molecules, the resulting MPF complex phosphorylates a variety of proteins, initiating mitosis. MPF acts both directly as a kinase and indirectly by activating other kinases. For example, MPF causes phosphorylation of various proteins of the nuclear lamina (see Figure 6.9), which promotes fragmentation of the nuclear envelope during prometaphase of mitosis. There is also evidence that MPF contributes to molecular events required for chromosome condensation and spindle formation during prophase.

During anaphase, MPF helps switch itself off by initiating a process that leads to the destruction of its own cyclin. The noncyclin part of MPF, the Cdk, persists in the cell, inactive until it becomes part of MPF again by associating with new cyclin molecules synthesized during the S and G_2 phases of the next round of the cycle.

Cell behavior at the G_1 checkpoint is also regulated by the activity of cyclin-Cdk protein complexes. Animal cells appear to have at least three Cdk proteins and several different cyclins that operate at this checkpoint. The fluctuating activities of different cyclin-Cdk complexes are of major importance in controlling all the stages of the cell cycle.

Stop and Go Signs: Internal and External Signals at the Checkpoints

Research scientists are currently working out the pathways that link signals originating inside and outside the cell with the responses by cyclin-dependent kinases and other proteins. An example of an internal signal occurs at the third important checkpoint, the M phase checkpoint. Anaphase, the separation of sister chromatids, does not begin until all the chromosomes are properly attached to the spindle at the metaphase plate. Researchers have learned that as long as some kinetochores are unattached to spindle microtubules, the sister chromatids remain together, delaying anaphase. Only when the kinetochores of all the chromosomes are properly attached to the spindle does the appropriate regulatory protein complex become activated. (In this case, the regulatory molecule is not a cyclin-Cdk complex but, instead, a different complex made up of several proteins.) Once activated, the complex sets off a chain of molecular events that activates the enzyme separase, which cleaves the cohesins, allowing the sister chromatids to separate. This mechanism ensures that daughter cells do not end up with missing or extra chromosomes.

Studies using animal cells in culture have led to the identification of many external factors, both chemical and physical, that can influence cell division. For example, cells fail to divide if an essential nutrient is lacking in the culture medium. (This is analogous to trying to run an automatic washing machine without the water supply hooked up; an internal sensor won't allow the machine to continue past the point where water is needed.) And even if all other conditions are favorable, most types of mammalian cells divide in culture only if the growth medium includes specific growth factors. As mentioned in Chapter 11, a **growth factor** is a protein released by certain cells that stimulates other cells to divide. Researchers have discovered more than 50 growth factors. Different cell types respond specifically to different growth factors or combinations of growth factors.



◀ **Figure 12.18 The effect of platelet-derived growth factor (PDGF) on cell division.**



(a) Normal mammalian cells. Contact with neighboring cells and the availability of nutrients, growth factors, and a substratum for attachment limit cell density to a single layer.



(b) Cancer cells. Cancer cells usually continue to divide well beyond a single layer, forming a clump of overlapping cells. They do not exhibit anchorage dependence or density-dependent inhibition.

▲ **Figure 12.19 Density-dependent inhibition and anchorage dependence of cell division.** Individual cells are shown disproportionately large in the drawings.

Consider, for example, *platelet-derived growth factor (PDGF)*, which is made by blood cell fragments called platelets. The experiment illustrated in **Figure 12.18** demonstrates that PDGF is required for the division of cultured fibroblasts, a type of connective tissue cell. Fibroblasts have PDGF receptors on their plasma membranes. The binding of PDGF molecules to these receptors (which are receptor tyrosine kinases; see Chapter 11) triggers a signal transduction pathway that allows the cells to pass the G_1 checkpoint and divide. PDGF stimulates fibroblast division not only in the artificial conditions of cell culture, but also in an animal's body. When an injury occurs, platelets release PDGF in the vicinity. The resulting proliferation of fibroblasts helps heal the wound.

The effect of an external physical factor on cell division is clearly seen in **density-dependent inhibition**, a phenomenon in which crowded cells stop dividing **(Figure 12.19a)**. As first observed many years ago, cultured cells normally divide until they form a single layer of cells on the inner surface of the culture container, at which point the cells stop dividing. If some cells are removed, those bordering the open space begin dividing again and continue until the vacancy is filled. Follow-up studies revealed that the binding of a cell-surface protein to its counterpart on an adjoining cell sends a growth-inhibiting signal to both cells, preventing them from moving forward in the cell cycle, even in the presence of growth factors.

Most animal cells also exhibit **anchorage dependence** (see Figure 12.19a). To divide, they must be attached to a substratum, such as the inside of a culture jar or the extracellular matrix of a tissue. Experiments suggest that like cell density,

anchorage is signaled to the cell cycle control system via pathways involving plasma membrane proteins and elements of the cytoskeleton linked to them.

Density-dependent inhibition and anchorage dependence appear to function in the body's tissues as well as in cell culture, checking the growth of cells at some optimal density and location. Cancer cells, which we discuss next, exhibit neither density-dependent inhibition nor anchorage dependence **(Figure 12.19b)**.

Loss of Cell Cycle Controls in Cancer Cells

Cancer cells do not heed the normal signals that regulate the cell cycle. They divide excessively and invade other tissues. If unchecked, they can kill the organism.

Cancer cells in culture do not stop dividing when growth factors are depleted. A logical hypothesis is that cancer cells do not need growth factors in their culture medium to grow and divide. They may make a required growth factor themselves, or they may have an abnormality in the signaling pathway that conveys the growth factor's signal to the cell cycle control system even in the absence of that factor. Another possibility is an abnormal cell cycle control system. In all of these scenarios, the underlying basis of the abnormality is almost always a change in one or more genes that alters the function of their protein products, resulting in faulty cell cycle control. You will learn more in Chapter 18 about the genetic bases of these changes and how these conditions may lead to cancer.

There are other important differences between normal cells and cancer cells that reflect derangements of the cell cycle. If and when they stop dividing, cancer cells do so at random points in the cycle, rather than at the normal checkpoints. Moreover, cancer cells can go on dividing indefinitely in culture if they are given a continual supply of nutrients; in essence, they are "immortal." A striking example is a cell line that has been reproducing in culture since 1951. Cells of this line are called HeLa cells because their original source was a tumor removed from a woman named Henrietta Lacks. By contrast, nearly all normal mammalian cells growing in culture divide only about 20 to 50 times before they stop dividing, age, and die. (We'll see a possible reason for this phenomenon when we discuss DNA replication in Chapter 16.) Finally, cancer cells evade the normal controls that trigger a cell to undergo apoptosis when something is wrong—for example, when an irreparable mistake has occurred during DNA replication preceding mitosis.

The abnormal behavior of cancer cells can be catastrophic when it occurs in the body. The problem begins when a single cell in a tissue undergoes **transformation**, the process that converts a normal cell to a cancer cell. The body's immune system normally recognizes a transformed cell as an insurgent and destroys it. However, if the cell evades destruction, it may proliferate and form a tumor, a mass of abnormal cells within otherwise normal tissue. The abnormal cells may remain at the original site if they have too few genetic and cellular changes to survive at another site. In that case, the tumor is called a **benign tumor**. Most benign tumors do not cause serious problems and can be completely removed by surgery. In contrast, a **malignant tumor** includes cells whose genetic and cellular changes enable them to spread to new tissues and impair the functions of one or more organs. An individual with a malignant tumor is said to have cancer; **Figure 12.20** shows the development of breast cancer.

The changes that have occurred in cells of malignant tumors show up in many ways besides excessive proliferation. These cells may have unusual numbers of chromosomes,



▲ **Figure 12.20 The growth and metastasis of a malignant breast tumor.** The cells of malignant (cancerous) tumors grow in an uncontrolled way and can spread to neighboring tissues and, via lymph and blood vessels, to other parts of the body. The spread of cancer cells beyond their original site is called metastasis.

though whether this is a cause or an effect of transformation is a current topic of debate. Their metabolism may be disabled, and they may cease to function in any constructive way. Abnormal changes on the cell surface cause cancer cells to lose attachments to neighboring cells and the extracellular matrix, allowing them to spread into nearby tissues. Cancer cells may also secrete signaling molecules that cause blood vessels to grow toward the tumor. A few tumor cells may separate from the original tumor, enter blood vessels and lymph vessels, and travel to other parts of the body. There, they may proliferate and form a new tumor. This spread of cancer cells to locations distant from their original site is called **metastasis** (see Figure 12.20).

A tumor that appears to be localized may be treated with high-energy radiation, which damages DNA in cancer cells much more than it does in normal cells, apparently because the majority of cancer cells have lost the ability to repair such damage. To treat known or suspected metastatic tumors, chemotherapy is used, in which drugs that are toxic to actively dividing cells are administered through the circulatory system. As you might expect, chemotherapeutic drugs interfere with specific steps in the cell cycle. For example, the drug Taxol freezes the mitotic spindle by preventing microtubule depolymerization, which stops actively dividing cells from proceeding past metaphase. The side effects of chemotherapy are due to the drugs' effects on normal cells that divide often. For example, nausea results from chemotherapy's effects on intestinal cells, hair loss from effects on hair follicle cells, and susceptibility to infection from effects on immune system cells.

Over the past several decades, researchers have produced a flood of valuable information about cell-signaling pathways and how their malfunction contributes to the development of cancer through effects on the cell cycle. Coupled with new molecular techniques, such as the ability to rapidly sequence the DNA of cells in a particular tumor, medical treatments for cancer are beginning to become more "personalized" to a particular patient's tumor. Breast cancer provides a good example. Basic research on the processes described in Chapters 11 and 12 has augmented our understanding of the molecular events underlying development of breast cancer. Proteins functioning in cell signaling pathways that affect the cell cycle are often found to be altered in breast cancer cells. Analyzing the level and sequences of such proteins has allowed physicians to better tailor the treatment to the cancers of some individuals, as shown in **Figure 12.21**.

One of the big lessons we've learned about the development of cancer, though, is how very complex the process is. There are many areas that remain to be explored. Perhaps the reason we have so many unanswered questions about cancer cells is that there is still so much to learn about how normal cells function. The cell, life's basic unit of structure and function, holds enough secrets to engage researchers well into the future.

▼ **Figure 12.21**

IMPACT

Advances in Treatment of Breast Cancer

Cancer cells, such as the breast cancer cell shown below, are analyzed by DNA sequencing and other molecular techniques to look for alterations in the level or sequence of specific proteins associated with cancer. For example, the cells of roughly 20–25% of breast cancer tumors show abnormally high amounts of a cell-surface receptor tyrosine kinase called HER2, and many show an increase in the number of estrogen receptor (ER) molecules, intracellular receptors that can trigger cell division. Based on lab findings, a physician can prescribe chemotherapy with a molecule that blocks the function of the specific protein (Herceptin for HER2 and tamoxifen for ERs). Treatment using these agents, when appropriate, has led to increased survival rates and fewer cancer recurrences.



WHY IT MATTERS Approximately one out of every eight women will develop breast cancer, the most common cancer among women. Worldwide, the incidence of breast cancer has been increasing annually. However, the mortality rate from this disease is falling in the United States and elsewhere, probably a result of earlier detection and improved treatment. Furthermore, what we are learning from the study of breast cancer also enhances our understanding of the development and treatment of other types of cancer.

FURTHER READING F. J. Esteva and G. N. Hortobagyi, Gaining ground on breast cancer, *Scientific American* 298:58–65 (2008).

MAKE CONNECTIONS Review the material in Chapter 11 on receptor tyrosine kinases and intracellular receptors (Figures 11.7 and 11.9 on pp. 212–214). Explain in general how these receptors might function in triggering cell division.

CONCEPT CHECK 12.3

1. In Figure 12.14, why do the nuclei resulting from experiment 2 contain different amounts of DNA?
2. How does MPF allow a cell to pass the G_2 phase checkpoint and enter mitosis? (See Figure 12.17.)
3. What phase are most of your body cells in?
4. Compare and contrast a benign tumor and a malignant tumor.
5. **WHAT IF?** What would happen if you performed the experiment in Figure 12.18 with cancer cells?

For suggested answers, see Appendix A.

12 CHAPTER REVIEW

SUMMARY OF KEY CONCEPTS

- Unicellular organisms reproduce by **cell division**; multicellular organisms depend on cell division for their development from a fertilized egg and for growth and repair. Cell division is part of the **cell cycle**, an ordered sequence of events in the life of a cell from its origin until it divides into daughter cells.

CONCEPT 12.1

Most cell division results in genetically identical daughter cells (pp. 229–230)

- The genetic material (DNA) of a cell—its **genome**—is partitioned among **chromosomes**. Each eukaryotic chromosome consists of one DNA molecule associated with many proteins that maintain chromosome structure and help control the activity of genes. Together, the complex of DNA and associated proteins is called **chromatin**. The chromatin of a chromosome exists in different states of condensation at different times. In animals, **gametes** have one set of chromosomes and **somatic cells** have two sets.
- Cells replicate their genetic material before they divide, ensuring that each daughter cell can receive a copy of the DNA. In preparation for cell division, chromosomes are duplicated, each one then consisting of two identical **sister chromatids** joined along their lengths by sister chromatid cohesion and held most tightly together at a constricted region at the **centromeres** of the chromatids. When this cohesion is broken, the chromatids separate during cell division, becoming the chromosomes of the new daughter cells. Eukaryotic cell division consists of **mitosis** (division of the nucleus) and **cytokinesis** (division of the cytoplasm).

? *Differentiate between these terms: chromosome, chromatin, and chromatid.*

CONCEPT 12.2

The mitotic phase alternates with interphase in the cell cycle (pp. 230–238)

- Between divisions, a cell is in **interphase**: the $\mathbf{G_1}$, **S**, and $\mathbf{G_2}$ phases. The cell grows throughout interphase, but DNA is replicated only during the synthesis (S) phase. Mitosis and cytokinesis make up the **mitotic (M) phase** of the cell cycle.



- The **mitotic spindle** is an apparatus of microtubules that controls chromosome movement during mitosis. In animal cells, the spindle arises from the **centrosomes** and includes spindle microtubules and **asters**. Some spindle microtubules attach to the **kinetochores** of chromosomes and move the chromosomes to the **metaphase plate**. In anaphase, sister chromatids separate, and motor proteins move them along the kinetochore microtubules toward opposite ends of the cell. Meanwhile, motor proteins push nonkinetochore microtubules from opposite poles away from each other, elongating the cell. In telophase, genetically identical daughter nuclei form at opposite ends of the cell.
- Mitosis is usually followed by cytokinesis. Animal cells carry out cytokinesis by **cleavage**, and plant cells form a **cell plate**.
- During **binary fission** in bacteria, the chromosome replicates and the two daughter chromosomes actively move apart. Some of the proteins involved in bacterial binary fission are related to eukaryotic actin and tubulin.
- Since prokaryotes preceded eukaryotes by more than a billion years, it is likely that mitosis evolved from prokaryotic cell division. Certain unicellular eukaryotes exhibit mechanisms of cell division that may be similar to those of ancestors of existing eukaryotes. Such mechanisms might have been intermediate steps in the evolution of mitosis from bacterial binary fission.

? *In which of the three subphases of interphase and the stages of mitosis do chromosomes exist as single DNA molecules?*

CONCEPT 12.3

The eukaryotic cell cycle is regulated by a molecular control system (pp. 238–243)

- Signaling molecules present in the cytoplasm regulate progress through the cell cycle.
- The **cell cycle control system** is molecularly based. Cyclic changes in regulatory proteins work as a cell cycle clock. The key molecules are **cyclins** and **cyclin-dependent kinases (Cdks)**. The clock has specific **checkpoints** where the cell cycle stops until a go-ahead signal is received. Cell culture has enabled researchers to study the molecular details of cell division. Both internal signals and external signals control the cell cycle checkpoints via signal transduction pathways. Most cells exhibit **density-dependent inhibition** of cell division as well as **anchorage dependence**.
- Cancer cells elude normal cell cycle regulation and divide out of control, forming tumors. **Malignant tumors** invade surrounding tissues and can undergo **metastasis**, exporting cancer cells to other parts of the body, where they may form secondary tumors. Recent advances in understanding the cell cycle and cell signaling, as well as techniques for sequencing DNA, have allowed improvements in cancer treatment.

? *Explain the significance of the G_1, G_2, and M checkpoints and the go-ahead signals involved in the cell cycle control system.*

TEST YOUR UNDERSTANDING

LEVEL 1: KNOWLEDGE/COMPREHENSION

1. Through a microscope, you can see a cell plate beginning to develop across the middle of a cell and nuclei forming on either side of the cell plate. This cell is most likely
   a. an animal cell in the process of cytokinesis.
   b. a plant cell in the process of cytokinesis.

c. an animal cell in the S phase of the cell cycle.
d. a bacterial cell dividing.
e. a plant cell in metaphase.

2. Vinblastine is a standard chemotherapeutic drug used to treat cancer. Because it interferes with the assembly of microtubules, its effectiveness must be related to
a. disruption of mitotic spindle formation.
b. inhibition of regulatory protein phosphorylation.
c. suppression of cyclin production.
d. myosin denaturation and inhibition of cleavage furrow formation.
e. inhibition of DNA synthesis.

3. One difference between cancer cells and normal cells is that cancer cells
a. are unable to synthesize DNA.
b. are arrested at the S phase of the cell cycle.
c. continue to divide even when they are tightly packed together.
d. cannot function properly because they are affected by density-dependent inhibition.
e. are always in the M phase of the cell cycle.

4. The decline of MPF activity at the end of mitosis is due to
a. the destruction of the protein kinase Cdk.
b. decreased synthesis of Cdk.
c. the degradation of cyclin.
d. the accumulation of cyclin.
e. synthesis of DNA.

5. In the cells of some organisms, mitosis occurs without cytokinesis. This will result in
a. cells with more than one nucleus.
b. cells that are unusually small.
c. cells lacking nuclei.
d. destruction of chromosomes.
e. cell cycles lacking an S phase.

6. Which of the following does *not* occur during mitosis?
a. condensation of the chromosomes
b. replication of the DNA
c. separation of sister chromatids
d. spindle formation
e. separation of the spindle poles

LEVEL 2: APPLICATION/ANALYSIS

7. In the light micrograph below of dividing cells near the tip of an onion root, identify a cell in each of the following stages: prophase, prometaphase, metaphase, anaphase, and telophase. Describe the major events occurring at each stage.



8. A particular cell has half as much DNA as some other cells in a mitotically active tissue. The cell in question is most likely in
a. G_1.
b. G_2.
c. prophase.
d. metaphase.
e. anaphase.

9. The drug cytochalasin B blocks the function of actin. Which of the following aspects of the animal cell cycle would be most disrupted by cytochalasin B?
a. spindle formation
b. spindle attachment to kinetochores
c. DNA synthesis
d. cell elongation during anaphase
e. cleavage furrow formation and cytokinesis

10. **DRAW IT** Draw one eukaryotic chromosome as it would appear during interphase, during each of the stages of mitosis, and during cytokinesis. Also draw and label the nuclear envelope and any microtubules attached to the chromosome(s).

LEVEL 3: SYNTHESIS/EVALUATION

11. **EVOLUTION CONNECTION**
The result of mitosis is that the daughter cells end up with the same number of chromosomes that the parent cell had. Another way to maintain the number of chromosomes would be to carry out cell division first and then duplicate the chromosomes in each daughter cell. Do you think this would be an equally good way of organizing the cell cycle? Why do you suppose that evolution has not led to this alternative?

12. **SCIENTIFIC INQUIRY**
Although both ends of a microtubule can gain or lose subunits, one end (called the plus end) polymerizes and depolymerizes at a higher rate than the other end (the minus end). For spindle microtubules, the plus ends are in the center of the spindle, and the minus ends are at the poles. Motor proteins that move along microtubules specialize in walking either toward the plus end or toward the minus end; the two types are called plus end–directed and minus end–directed motor proteins, respectively. Given what you know about chromosome movement and spindle changes during anaphase, predict which type of motor proteins would be present on (a) kinetochore microtubules and (b) nonkinetochore microtubules.

13. **WRITE ABOUT A THEME**
The Genetic Basis of Life The continuity of life is based on heritable information in the form of DNA. In a short essay (100–150 words), explain how the process of mitosis faithfully parcels out exact copies of this heritable information in the production of genetically identical daughter cells.

For selected answers, see Appendix A.

MasteringBIOLOGY® www.masteringbiology.com

1. MasteringBiology® Assignments

BioFlix **Tutorials** Mitosis: Mitosis and the Cell Cycle • Mechanism of Mitosis • Comparing Cell Division in Animals, Plants, and Bacteria
Activities The Cell Cycle • Mitosis and Cytokinesis Animation • Four Phases of the Cell Cycle • Causes of Cancer • Discovery Channel Video: Fighting Cancer
Questions Student Misconceptions • Reading Quiz • Multiple Choice • End-of-Chapter

2. eText

Read your book online, search, take notes, highlight text, and more.

3. The Study Area

Practice Tests • Cumulative Test • BioFlix 3-D Animations • MP3 Tutor Sessions • Videos · Activities • Investigations • Lab Media • Audio Glossary • Word Study Tools • Art